

Sheet 5**Exercise 5.1**

Let \mathcal{H} be a Hilbert space, $f \in \mathcal{H}$ and define a linear functional by

$$\Phi(f) : \mathcal{H} \rightarrow \mathbb{C}, \quad g \mapsto \langle f, g \rangle.$$

Show that

$$\|\Phi(f)\| = \|f\|.$$

Exercise 5.2

The adjoint has the following properties for $A, B \in \mathcal{B}(\mathcal{H})$ and $z \in \mathbb{C}$

- a) $(A + zB)^* = A^* + \bar{z}B^*$;
- b) $(AB)^* = B^*A^*$;
- c) $(A^*)^* = A$
- d) $\ker A^* = (\text{ran } A)^\perp$ and $\ker A = (\text{ran } A^*)^\perp$.

Homework

Exercise 5.3 (The Lax-Milgram Theorem)

Let \mathcal{H} be a Hilbert space and

$$\alpha : \mathcal{H} \times \mathcal{H} \rightarrow \mathbb{C}$$

a sesquilinear form. Assume that

- α is *bounded*: there exists $C > 0$ so that for all $f, g \in \mathcal{H}$

$$|\alpha(f, g)| \leq C \|f\| \|g\|;$$

- α is *coercive*: there exists $a > 0$ so that for all $f \in \mathcal{H}$

$$\alpha(f, f) \geq a \|f\|^2.$$

Prove that:

- There exists $A \in \mathcal{B}(\mathcal{H})$ so that $\alpha(f, g) = \langle Af, g \rangle$;
- A is bijective with bounded inverse satisfying $\|A^{-1}\| \leq a^{-1}$;
- $g = A^{-1}f$ is the unique minimiser of

$$g \mapsto \alpha(g, g) - 2\operatorname{Re}\langle f, g \rangle.$$

Hand in on 08.03.2023 before the lecture or by 10:00 by mail to jonas.lampart@u-bourgogne.fr.